


**PCT**WORLD INTELLECTUAL PROPERTY  
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER T.

WO 9606087A1

(51) International Patent Classification <sup>6</sup> : <b>C07D 243/14, 223/16, 267/14, 281/10, A61K 31/55</b>		A1	(11) International Publication Number: <b>WO 96/06087</b> (43) International Publication Date: 29 February 1996 (29.02.96)
(21) International Application Number: PCT/US95/10670 (22) International Filing Date: 22 August 1995 (22.08.95)  (30) Priority Data: 08/294,278 22 August 1994 (22.08.94) US  (60) Parent Application or Grant (63) Related by Continuation US 08/294,278 (CON) Filed on 22 August 1994 (22.08.94)  (71) Applicant (for all designated States except US): SMITHKLINE BEECHAM CORPORATION [US/US]; Corporate Intellectual Property, UW 2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US).  (72) Inventors; and (75) Inventors/Applicants (for US only): KWON, Chet [US/US]; Apartment D503, 649 South Henderson Road, King of Prussia, PA 19406 (US). MILLER, William, Henry [US/US]; 333 Fell Lane, Schwenksville, PA 19473 (US).			(74) Agents: McCARTHY, Mary, E. et al.; SmithKline Beecham Corporation, Corporate Intellectual Property, UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US).  (81) Designated States: JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  Published With international search report.
(54) Title: BICYCLIC COMPOUNDS			
(57) Abstract <p>This invention relates to compounds of formula (I) or a pharmaceutically acceptable salt thereof, which are effective for inhibiting platelet aggregation and bone resorption, pharmaceutical compositions for effecting such activity, and a method for using these compounds.</p>			



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgystan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

## Bicyclic Compounds

### Field of the Invention

This invention relates to pharmaceutically active bicyclic compounds,  
5 pharmaceutical compositions containing the compounds and methods of using the  
compounds. The instant compounds inhibit platelet aggregation. These compounds  
also inhibit the vitronectin receptor and are useful for the treatment of osteoporosis.

### Background of the Invention

10 Platelet aggregation is believed to be mediated primarily through the  
fibrinogen receptor, or GPIIb-IIIa platelet receptor complex, which is a member of a  
family of adhesion receptors referred to as integrins. It has been found that  
frequently the natural ligands of integrin receptors are proteins which contain an  
Arg-Gly-Asp sequence. Von Willebrand factor and fibrinogen, which are  
15 considered to be natural ligands for the GPIIb-IIIa receptor, possess an Arg-Gly-Asp  
(RGD in single letter amino acid code) sequence in their primary structure.  
Functionally, these proteins are able to bind and crosslink GPIIb-IIIa receptors on  
adjacent platelets and thereby effect aggregation of platelets.

Fibronectin, vitronectin and thrombospondin are RGD-containing proteins  
20 which have also been demonstrated to bind to GPIIb-IIIa. Fibronectin is found in  
plasma and as a structural protein in the intracellular matrix. Binding between the  
structural proteins and GPIIb-IIIa may function to cause platelets to adhere to  
damaged vessel walls.

Also, recent studies have indicated that the attachment of osteoclasts to the  
25 bone matrix is mediated through cell surface adhesion receptors called integrins.  
For instance, Davies, *et al.*, *J. Cell Biol.* 1989, 109, 1817, disclose that the  
osteoclast functional antigen, which is implicated in the regulation of bone  
resorption, is biochemically related to the vitronectin receptor. The vitronectin  
receptor, or the  $\alpha_v\beta_3$  integrin, is known to bind to bone matrix proteins, such as  
30 osteopontin, bone sialoprotein and thrombospondin, which contain the tri-peptide  
Arg-Gly-Asp (or RGD) motif. Thus, Horton, *et al.*, *Exp. Cell Res.* 1991, 195, 368,  
disclose that RGD-containing peptides and an anti-vitronectin receptor antibody  
(23C6) inhibit dentine resorption and cell spreading by osteoclasts. In addition,  
Sato, *et al.*, *J. Cell Biol.* 1990, 111, 1713 disclose that echistatin, a snake venom  
35 peptide which contains the RGD sequence, is a potent inhibitor of bone resorption in  
tissue culture, and inhibits attachment of osteoclasts to bone. Fisher, *et al.*,  
*Endocrinology* 1993, 132, 1411, has further shown that echistatin inhibits bone

resorption *in vivo* in the rat. EP 528 587 and 528 586 report substituted phenyl derivatives which inhibit osteoclast mediated bone resorption.

The present invention discloses novel bicyclic compounds, including benzazepines and benzodiazepines, which are inhibitors of the GPIIb-IIIa receptor and inhibit platelet aggregation. Also, the instant compounds are inhibitors of the vitronectin receptor. These agents inhibit bone resorption and are useful for the treatment of osteoporosis.

#### Summary of the Invention

10 In one aspect this invention is a bicyclic compound comprising a substituted six-membered ring fused to a substituted seven-membered ring as described hereinafter in formula (I).

This invention is also a pharmaceutical composition which comprises a compound of formula (I) and a pharmaceutically acceptable carrier.

15 This invention is further a method for inhibiting platelet aggregation in a mammal in need thereof, which comprises internally administering an effective amount of a compound of formula (I).


In another aspect, this invention provides a method for inhibiting reocclusion of an artery or vein in a mammal following fibrinolytic therapy, which comprises internally administering an effective amount of a fibrinolytic agent and a compound of formula (I). This invention is also a method for treating stroke, transient ischemia attacks, or myocardial infarction.

20 This invention is also a method of treating diseases which are mediated by ligands which bind to the vitronectin receptor. In a particular aspect, the compounds of this invention are useful for treating osteoporosis and diseases wherein bone resorption is a factor.

#### Detailed Description of the Invention

30 This invention discloses bicyclic compounds which inhibit platelet aggregation and which inhibit bone resorption. The novel bicyclic compounds comprise a seven-membered ring fused to an aromatic six membered ring and having a nitrogen-containing substituent on the six membered ring and an aliphatic substituent containing an acidic moiety on the seven membered ring. The seven membered ring contains heteroatoms, such as nitrogen, oxygen and sulfur, and the six membered ring is carbocyclic.

The compounds of this invention are compounds of formula (I):



wherein:

- 5           A is NR', CHR', O or S;  
             X<sup>1</sup> and X<sup>2</sup> are C=O or CH<sub>2</sub>, with the proviso that only one of X<sup>1</sup> or X<sup>2</sup> is C=O;  
             R<sup>1</sup> is H, C<sub>1-6</sub>alkyl or (CHR')<sub>m</sub>-Y;  
             R<sup>2</sup> is -(CH<sub>2</sub>)<sub>t</sub>-CO<sub>2</sub>R<sup>3</sup>;  
 10          R<sup>3</sup> is H, C<sub>1-6</sub>alkyl or (CHR')<sub>m</sub>-Ar;  
             R<sup>4</sup> and R<sup>4</sup>' independently are -(CH<sub>2</sub>)<sub>s</sub>- $\textcircled{\text{N}}$  or -(CH<sub>2</sub>)<sub>s</sub>-NR'R'  
             Y is H, Ar, C<sub>3-7</sub>cycloalkyl, CO<sub>2</sub>R<sup>3</sup> or Tet;  
             R' is H, C<sub>1-6</sub>alkyl or (CH<sub>2</sub>)<sub>m</sub>-Ar;  
             Ar is phenyl or naphthyl unsubstituted or substituted by one to three  
 15          C<sub>1-6</sub>alkyl, trifluoromethyl, halogen, OR', SR', CONR'R<sup>1</sup>, CO<sub>2</sub>R<sup>1</sup>, NO<sub>2</sub> or R<sup>5</sup>R<sup>1</sup>N;  
             R<sup>5</sup> is H, C<sub>1-6</sub>alkyl, (CHR')<sub>m</sub>-Y or CO(CHR')<sub>m</sub>-Y;  
              $\textcircled{\text{N}}$  is pyrroline, pyrrolidine, imidazole, imidazoline, imidazolidine, pyrazole,  
             pyrazoline, pyrazolidine, piperidine, piperazine, morpholine, pyridine, pyridinium,  
             tetrahydropyridine, tetrahydroazepine or hexahydroazepine;  
 20          m is 0 to 6;  
             s is 1 to 4; and  
             t is 1 or 2;  
             or a pharmaceutically acceptable salt thereof.

- 25           Also included in this invention are pharmaceutically acceptable addition salts, complexes or prodrugs of the compounds of this invention. Prodrugs are considered to be any covalently bonded carriers which release the active parent drug according to formula (I) *in vivo*.

- 30           In cases wherein the compounds of this invention may have one or more chiral centers, unless specified, this invention includes each unique nonracemic compound which may be synthesized and resolved by conventional techniques.

The meaning of any substituent at any one occurrence is independent of its meaning, or any other substituent's meaning, at any other occurrence, unless specified otherwise.

With reference to formula (I), suitably,  $X^1$  is  $CH_2$ ,  $X^2$  is  $C=O$  and A is  $NR'$ .

5 Preferably, with reference to formula (I),

$X^1$  is  $CH_2$ ,  $X^2$  is  $C=O$ , A is  $NR'$ ,  $R^1$  is  $C_{1-4}$ alkyl or  $(CHR')_n-Y$ , in which  $R'$  is H or  $C_{1-4}$ alkyl, m is 1 to 3 and Y is Ar or  $C_{3-7}$ cycloalkyl, and  $R^4$  and  $R^4'$  independently are  $-(CH_2)_s-\textcircled{N}$ , in which  $\textcircled{N}$  is piperidine, piperazine, pyridine, pyridinium or tetrahydropyridine and s is 1 to 3. Most preferably,  $\textcircled{N}$  is pyridine or

10 piperidine.

Preferred compounds of this inventions are:

( $\pm$ )-7-[[bis[2-(4-pyridinyl)ethyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid,

( $\pm$ )-8-[[bis[2-(4-pyridinyl)ethyl]amino]carbonyl]-3-oxo-4-(2-phenylethyl)-  
15 2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid,

( $\pm$ )-7-[[bis[2-(4-piperidinyl)ethyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid, and

( $\pm$ )-8-[[bis[2-(4-piperidinyl)ethyl]amino]carbonyl]-4-(2-cyclohexylethyl)-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;

20 or a pharmaceutically acceptable salt thereof.

The most preferred compounds of this invention are:

( $\pm$ )-7-[[bis[2-(4-piperidinyl)ethyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid, and

( $\pm$ )-8-[[bis[2-(4-piperidinyl)ethyl]amino]carbonyl]-4-(2-cyclohexylethyl)-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;

or a pharmaceutically acceptable salt thereof.

In the above description,  $C_{1-4}$ alkyl is meant to include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl and t-butyl.  $C_{1-6}$ alkyl additionally includes  
30 pentyl, n-pentyl, isopentyl, neopentyl and hexyl and the simple aliphatic isomers thereof.

$C_{3-7}$ cycloalkyl refers to an optionally substituted carbocyclic system of three to seven carbon atoms, which may contain up to two unsaturated carbon-carbon bonds. Typical of  $C_{3-7}$ cycloalkyl are cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl and cycloheptyl, particularly cyclohexyl.  
35 Any combination of up to three substituents, such as chosen from  $C_{1-6}$ alkyl, trifluoromethyl, halogen,  $OR'$ ,  $SR'$ ,  $CONR'R^1$ ,  $CO_2R^1$ ,  $NO_2$  or  $R^5R^1N$ , on the




cycloalkyl ring that is available by conventional chemical synthesis and is stable, is within the scope of this invention.

The accessible substituted seven-membered ring as referred of formula (I) is any saturated seven-membered ring which (i) contains up to two heteroatoms  
5 selected from the group of N, O and S, wherein S and N may optionally be oxidized, and (ii) is stable and may be synthesized by one skilled in the chemical arts in a form fused via two adjacent ring carbon atoms to a phenyl. Typical of accessible seven-membered rings are the common saturated rings of azepine, diazepine, thiazepin and oxazepin. Representative bicyclic rings formed by the  
10 combination of the accessible seven-membered and phenyl rings are: benzaepine, benzodiazepine, benzoxazepine, and benzothiazepine compounds. Benzodiazepine compounds are the preferred bicyclic compounds of the instant invention.

Certain radical groups are abbreviated herein. t-Bu refers to the tertiary butyl radical, Boc refers to the t-butyloxycarbonyl radical, Fmoc refers to the  
15 fluorenylmethoxycarbonyl radical, Ph refers to the phenyl radical, Cbz refers to the benzyloxycarbonyl radical, BrZ refers to the o-bromobenzyloxycarbonyl radical, ClZ refers to the o-chlorobenzyloxycarbonyl radical, Bzl refers to the benzyl radical, 4-MBzl refers to the 4-methyl benzyl radical, Me refers to methyl, Et refers to ethyl, Ac refers to acetyl, Alk refers to C<sub>1-6</sub>alkyl, Nph refers to 1- or 2-naphthyl  
20 and cHex refers to cyclohexyl. MeArg is N<sup>α</sup>-methyl arginine. Tet refers to 5-tetrazolyl.

Certain reagents are abbreviated herein. DCC refers to dicyclohexylcarbodiimide, DMAP refers to dimethylaminopyridine, DIEA refers to diisopropylethyl amine, EDC refers to N-ethyl-N'(dimethylaminopropyl)-  
25 carbodiimide. HOBt refers to 1-hydroxybenzotriazole, THF refers to tetrahydrofuran, DMF refers to dimethyl formamide, NBS refers to N-bromosuccinimide, Pd/C refers to a palladium on carbon catalyst, PPA refers to 1-propanephosphonic acid cyclic anhydride, DPPA refers to diphenylphosphoryl azide, BOP refers to benzotriazol-1-yloxy-tris(dimethylamino)phosphonium  
30 hexafluorophosphate, HF refers to hydrofluoric acid, TEA refers to triethylamine, TFA refers to trifluoroacetic acid, PCC refers to pyridinium chlorochromate.

The compounds of formula (I) are generally prepared by reacting a compound of the formula (II) with a compound of the formula (III):



(II)



(III)


- 5 wherein  $X^1$ ,  $X^2$ ,  $R^1$ ,  $R^2$ , A and  $R^4$  and  $R^{4'}$  are as defined in formula (I), with any reactive functional groups protected; and thereafter removing the protecting groups, and optionally forming a pharmaceutically acceptable salt.

- 10 Compounds of Formula (II) are benzodiazepines, benzazepines, benzozepines, and benzothiazepines described in Bondinell, et al., PCT Publication No. WO 93/00095, (PCT/US92/05463) published January 7, 1993. Reference should be made to such published patent application for its full disclosure, which is incorporated herein by reference.

- 15 Compounds of the formula (I) are prepared by the general methods described in Schemes I and II.




## Scheme I




5

1



2

10



3


- 15 a) EDC, HOBT, (i-Pr)<sub>2</sub>NEt, DMF, bis[2-(4-pyridyl)ethyl]amine; b) 1.0 N LiOH, aqueous THF; c) acidification.

20 Methyl (±)-8-carboxy-3-oxo-4-(2-phenylethyl)-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (I-1), prepared as described by Bondinell, et al. (WO 93/00095), is converted to an activated form of the carboxylic acid using, for example, EDC and HOBT, or SOCl<sub>2</sub>, and the activated form is subsequently reacted with an appropriate triamine, such as bis [2-(4-pyridyl)ethyl]amine, to afford the corresponding amide I-2. Many additional methods for converting a carboxylic acid


into an amide are known, and can be found in standard reference books, such as "Compendium of Organic Synthetic Methods", Vol. I - VI (published by Wiley-Interscience). The methyl ester of I-2 is hydrolyzed using aqueous base, for example, aqueous LiOH in THF, or aqueous NaOH in methanol, and the  
5 intermediate carboxylate salt is acidified with a suitable acid, for instance TFA or HCl, to afford the carboxylic acid I-3. Alternatively, the intermediate carboxylate salt can be isolated, if desired.

## Scheme II


10



15



20



a) H<sub>2</sub>, PtO<sub>2</sub>, 1.0 N HCl, MeOH; b) see Scheme I.

Piperidine-containing compounds, such as II-3, can be prepared either from  
5 a suitably N-protected piperidine derivative, according to the methods described in  
Scheme 1, or from a pyridine precursor, such as II-1. For example, the pyridine  
subunits of II-1 can be reduced to the corresponding piperidine groups by  
hydrogenation over a suitable catalyst, preferably PtO<sub>2</sub>, in the presence of acid, such  
as HCl. Under these conditions, other aromatic rings in the molecule, such as may  
10 be present in the substituent at the 4-position, may be reduced simultaneously. The  
resulting piperidinium salt II-2 is then converted to compound II-3 by the methods  
described in Scheme I.

Coupling reagents as used herein denote reagents which may be used to form  
15 peptide bonds. Typical coupling methods employ carbodiimides, activated  
anhydrides and esters and acyl halides. Reagents such as EDC, DCC, DPPA, PPA,  
BOP reagent, HOBt, N-hydroxysuccinimide and oxalyl chloride are typical.

Coupling methods to form peptide bonds are generally well known to the art.  
The methods of peptide synthesis generally set forth by Bodansky *et al.*, THE  
20 PRACTICE OF PEPTIDE SYNTHESIS, Springer-Verlag, Berlin, 1984, Ali *et al.* in  
*J. Med. Chem.*, 29, 984 (1986) and *J. Med. Chem.*, 30, 2291 (1987) are generally  
illustrative of the technique and are incorporated herein by reference.

Solution synthesis for the formation of amide or peptide bonds is  
accomplished using conventional methods used to form amide bonds. Typically, the  
25 free amino group of a formula (III) HNR<sup>4</sup>R<sup>4'</sup> compound is coupled to an  
appropriate carboxylic acid substrate, such as a Scheme I, formula 1 compound,  
using a suitable carbodiimide coupling agent, such as N,N' dicyclohexyl  
carbodiimide (DCC), optionally in the presence of catalysts such as 1-  
hydroxybenzotriazole (HOBt) and dimethylamino pyridine (DMAP). Other  
30 methods, such as the formation of activated esters, anhydrides or acid halides, of the  
free carboxyl of, for example, a Scheme I, formula 1 compound, and subsequent  
reaction with the free amino group of a formula (III) HNR<sup>4</sup>R<sup>4'</sup> compound.

Acid addition salts of the compounds are prepared in a standard manner in a  
suitable solvent from the parent compound and an excess of an acid, such as  
35 hydrochloric, hydrobromic, sulfuric, phosphoric, acetic, maleic, succinic or  
methanesulfonic. Certain of the compounds form inner salts or zwitterions which  
may be acceptable. Cationic salts are prepared by treating the parent compound

with an excess of an alkaline reagent, such as a hydroxide, carbonate or alkoxide, containing the appropriate cation; or with an appropriate organic amine. Cations such as Li<sup>+</sup>, Na<sup>+</sup>, K<sup>+</sup>, Ca<sup>++</sup>, Mg<sup>++</sup> and NH<sub>4</sub><sup>+</sup> are specific examples of cations present in pharmaceutically acceptable salts.

5           This invention also provides a pharmaceutical composition which comprises a compound according to formula (I) and a pharmaceutically acceptable carrier. Accordingly, the compounds of formula (I) may be used in the manufacture of a medicament. Pharmaceutical compositions of the compounds of formula (I) prepared as hereinbefore described may be formulated as solutions or lyophilized  
10           powders for parenteral administration. Powders may be reconstituted by addition of a suitable diluent or other pharmaceutically acceptable carrier prior to use. The liquid formulation may be a buffered, isotonic, aqueous solution. Examples of suitable diluents are normal isotonic saline solution, standard 5% dextrose in water or buffered sodium or ammonium acetate solution. Such formulation is especially  
15           suitable for parenteral administration, but may also be used for oral administration or contained in a metered dose inhaler or nebulizer for insufflation. It may be desirable to add excipients such as polyvinylpyrrolidone, gelatin, hydroxy cellulose, acacia, polyethylene glycol, mannitol, sodium chloride or sodium citrate.

          Alternately, these compounds may be encapsulated, tableted or prepared in a  
20           emulsion or syrup for oral administration. Pharmaceutically acceptable solid or liquid carriers may be added to enhance or stabilize the composition, or to facilitate preparation of the composition. Solid carriers include starch, lactose, calcium sulfate dihydrate, terra alba, magnesium stearate or stearic acid, talc, pectin, acacia, agar or gelatin. Liquid carriers include syrup, peanut oil, olive oil, saline and water.  
25           The carrier may also include a sustained release material such as glyceryl monostearate or glyceryl distearate, alone or with a wax. The amount of solid carrier varies but, preferably, will be between about 20 mg to about 1 g per dosage unit. The pharmaceutical preparations are made following the conventional  
30           techniques of pharmacy involving milling, mixing, granulating, and compressing, when necessary, for tablet forms; or milling, mixing and filling for hard gelatin capsule forms. When a liquid carrier is used, the preparation will be in the form of a syrup, elixir, emulsion or an aqueous or non-aqueous suspension. Such a liquid formulation may be administered directly p.o. or filled into a soft gelatin capsule.

          For rectal administration, the compounds of this invention may also be  
35           combined with excipients such as cocoa butter, glycerin, gelatin or polyethylene glycols and molded into a suppository.

The compounds of this invention may be used *in vitro* to inhibit the aggregation of platelets in blood and blood products, *e.g.*, for storage, or for *ex vivo* manipulations such as in diagnostic or research use.

This invention also provides a method of inhibiting platelet aggregation and clot formation in a mammal, especially a human, which comprises the internal administration of a compound of formula (I) and a pharmaceutically acceptable carrier. Indications for such therapy include acute myocardial infarction (AMI), deep vein thrombosis, pulmonary embolism, dissecting aneurysm, transient ischemia attack (TIA), stroke and other infarct-related disorders, and unstable angina.

Chronic or acute states of hyper-aggregability, such as disseminated intravascular coagulation (DIC), septicemia, surgical or infectious shock, post-operative and post-partum trauma, cardiopulmonary bypass surgery, incompatible blood transfusion, abruptio placenta, thrombotic thrombocytopenic purpura (TTP), snake venom and immune diseases, are likely to be responsive to such treatment. In addition, the compounds of this invention may be useful in a method for the prevention of metastatic conditions, the prevention or treatment of fungal or bacterial infection, inducing immunostimulation and the treatment of sickle cell disease.

The compounds described herein are antagonists of the vitronectin receptor, and are useful for treating diseases wherein the underlying pathology is attributable to ligand or cell which interacts with the vitronectin receptor. For instance, these compounds are useful for the treatment of diseases wherein loss of the bone matrix creates pathology. Thus, the instant compounds are useful for the treatment of osteoporosis, hyperparathyroidism, Paget's disease, hypercalcemia of malignancy, osteolytic lesions produced by bone metastasis, bone loss due to immobilization or sex hormone deficiency. The compounds of this invention are also believed to have utility as antitumor and antiinflammatory agents, and be useful in the treatment of atherosclerosis and restenosis.

The compound is administered either orally or parenterally to the patient, in a manner such that the concentration of drug in the plasma is sufficient to inhibit platelet aggregation or bone resorption, or other such indication. The pharmaceutical composition containing the compound is administered at a dose between about 0.2 to about 50 mg/kg in a manner consistent with the condition of the patient. For acute therapy, parenteral administration is preferred. An intravenous infusion of the compound in 5% dextrose in water or normal saline, or a similar formulation with suitable excipients, is most effective, although an intramuscular bolus injection is also useful. Typically, the parenteral dose will be about 0.01 to about 100 mg/kg; preferably between 0.1 to 20 mg/kg. The

compounds are administered one to four times daily at a level to achieve a total daily dose of about 0.4 to about 400 mg/kg/day. The precise level and method by which the compounds are administered is readily determined by one skilled in the art by comparing the blood level of the agent to the concentration required to have a therapeutic effect.

This invention further provides a method for inhibiting the reocclusion of an artery or vein following fibrinolytic therapy, which comprises internal administration of a peptide of formula (I) and a fibrinolytic agent. It has been found that administration of a peptide in fibrinolytic therapy either prevents reocclusion completely or prolongs the time to reocclusion.

When used in the context of this invention the term fibrinolytic agent is intended to mean any compound, whether a natural or synthetic product, which directly or indirectly causes the lysis of a fibrin clot. Plasminogen activators are a well known group of fibrinolytic agents. Useful plasminogen activators include, for example, anistreplase, urokinase (UK), pro-urokinase (pUK), streptokinase (SK), tissue plasminogen activator (tPA) and mutants, or variants, thereof, which retain plasminogen activator activity, such as variants which have been chemically modified or in which one or more amino acids have been added, deleted or substituted or in which one or more functional domains have been added, deleted or altered such as by combining the active site of one plasminogen activator with the fibrin binding domain of another plasminogen activator or fibrin binding molecule. Other illustrative variants include tPA molecules in which one or more glycosylation sites have been altered. Preferred among plasminogen activators are variants of tPA in which the primary amino acid sequence has been altered in the growth factor domain so as to increase the serum half-life of the plasminogen activator. tPA Growth factor variants are disclosed, *e.g.*, by Robinson *et al.*, EP-A 0 297 589 and Browne *et al.*, EP-A 0 240 334. Other variants include hybrid proteins, such as those disclosed in EP 0 028 489, EP 0 155 387 and EP 0 297 882, all of which are incorporated herein by reference. Anistreplase is a preferred hybrid protein for use in this invention. Fibrinolytic agents may be isolated from natural sources, but are commonly produced by traditional methods of genetic engineering.

Useful formulations of tPA, SK, UK and pUK are disclosed, for example, in EP-A 0 211 592, EP-A 0 092 182 and U.S. Patent 4,568,543, all of which are incorporated herein by reference. Typically the fibrinolytic agent may be formulated in an aqueous, buffered, isotonic solution, such as sodium or ammonium acetate or adipate buffered at pH 3.5 to 5.5. Additional excipients such as polyvinyl



pyrrolidone, gelatin, hydroxy cellulose, acacia, polyethylene, glycol, mannitol and sodium chloride may also be added. Such a composition can be lyophilized.

The pharmaceutical composition may be formulated with both the compound of formula (I) and fibrinolytic in the same container, but formulation in different  
5 containers is preferred. When both agents are provided in solution form they can be contained in an infusion/injection system for simultaneous administration or in a tandem arrangement.

Indications for such therapy include myocardial infarction, deep vein thrombosis, pulmonary embolism, stroke and other infarct-related disorders. The  
10 peptide is administered just prior to, at the same time as, or just after parenteral administration of tPA or other fibrinolytic agent. It may prove desirable to continue treatment with the peptide for a period of time well after reperfusion has been established to maximally inhibit post-therapy reocclusion. The effective dose of tPA, SK, UK or pUK may be from 0.5 to 5 mg/kg and the effective dose of the  
15 peptide may be from about 0.1 to 25 mg/kg.

For convenient administration of the inhibitor and the fibrinolytic agent at the same or different times, a kit is prepared, comprising, in a single container, such as a box, carton or other container, individual bottles, bags, vials or other containers each having an effective amount of the inhibitor for parenteral administration, as  
20 described above, and an effective amount of tPA, or other fibrinolytic agent, for parenteral administration, as described above. Such kit can comprise, for example, both pharmaceutical agents in separate containers or the same container, optionally as lyophilized plugs, and containers of solutions for reconstitution. A variation of this is to include the solution for reconstitution and the lyophilized plug in two  
25 chambers of a single container, which can be caused to admix prior to use. With such an arrangement, the fibrinolytic and the peptide may be packaged separately, as in two containers, or lyophilized together as a powder and provided in a single container.

When both agents are provided in solution form, they can be contained in an  
30 infusion/injection system for simultaneous administration or in a tandem arrangement. For example, the platelet aggregation inhibitor may be in an i.v. injectable form, or infusion bag linked in series, via tubing, to the fibrinolytic agent in a second infusion bag. Using such a system, a patient can receive an initial bolus-type injection or infusion, of the peptide inhibitor followed by an infusion of the  
35 fibrinolytic agent.



The compounds may be tested in one of several biological assays to determine the concentration of compound which is required to have a given pharmacological effect.

## 5                                    Inhibition of RGD-mediated GPIIb-IIIa binding

### Purification of GPIIb-IIIa

Ten units of outdated, washed human platelets (obtained from Red Cross) were lyzed by gentle stirring in 3% octylglucoside, 20 mM Tris-HCl, pH 7.4, 140  
10 mM NaCl, 2 mM CaCl<sub>2</sub> at 4°C for 2 h. The lysate was centrifuged at 100,000g for 1 h. The supernatant obtained was applied to a 5 mL lentil lectin sepharose 4B column (E.Y. Labs) preequilibrated with 20 mM Tris-HCl, pH 7.4, 100 mM NaCl, 2 mM CaCl<sub>2</sub>, 1% octylglucoside (buffer A). After 2 h incubation, the column was washed with 50 mL cold buffer A. The lectin-retained GPIIb-IIIa was eluted with  
15 buffer A containing 10% dextrose. All procedures were performed at 4°C. The GPIIb-IIIa obtained was >95% pure as shown by SDS polyacrylamide gel electrophoresis.

### Incorporation of GPIIb-IIIa in Liposomes.

20                    A mixture of phosphatidylserine (70%) and phosphatidylcholine (30%) (Avanti Polar Lipids) were dried to the walls of a glass tube under a stream of nitrogen. Purified GPIIb-IIIa was diluted to a final concentration of 0.5 mg/mL and mixed with the phospholipids in a protein:phospholipid ratio of 1:3 (w:w). The mixture was resuspended and sonicated in a bath sonicator for 5 min. The mixture  
25 was then dialyzed overnight using 12,000-14,000 molecular weight cutoff dialysis tubing against a 1000-fold excess of 50 mM Tris-HCl, pH 7.4, 100 mM NaCl, 2 mM CaCl<sub>2</sub> (with 2 changes). The GPIIb-IIIa-containing liposomes were centrifuged at 12,000g for 15 min and resuspended in the dialysis buffer at a final protein concentration of approximately 1 mg/mL. The liposomes were stored at -70°C until  
30 needed.

### Competitive Binding to GPIIb-IIIa

The binding to the fibrinogen receptor (GPIIb-IIIa) was assayed by an indirect competitive binding method using [<sup>3</sup>H]-SK&F-107260 as an RGD-type  
35 ligand. The binding assay was performed in a 96-well filtration plate assembly (Millipore Corporation, Bedford, MA) using 0.22 µm hydrophilic durapore membranes. The wells were precoated with 0.2 mL of 10 µg/mL polylysine (Sigma

Chemical Co., St. Louis, MO.) at room temperature for 1 h to block nonspecific binding. Various concentrations of unlabeled benzodiazapines were added to the wells in quadruplicate. [<sup>3</sup>H]-SK&F-107260 was applied to each well at a final concentration of 4.5 nM, followed by the addition of 1 µg of the purified platelet GPIIb-IIIa-containing liposomes. The mixtures were incubated for 1 h at room temperature. The GPIIb-IIIa-bound [<sup>3</sup>H]-SK&F-107260 was separated from the unbound by filtration using a Millipore filtration manifold, followed by washing with ice-cold buffer (2 times, each 0.2 mL). Bound radioactivity remaining on the filters was counted in 1.5 mL Ready Solve (Beckman Instruments, Fullerton, CA) in a Beckman Liquid Scintillation Counter (Model LS6800), with 40% efficiency. Nonspecific binding was determined in the presence of 2 µM unlabeled SK&F-107260 and was consistently less than 0.14% of the total radioactivity added to the samples. All data points are the mean of quadruplicate determinations.

Competition binding data were analyzed by a nonlinear least-squares curve fitting procedure. This method provides the IC<sub>50</sub> of the antagonists (concentration of the antagonist which inhibits specific binding of [<sup>3</sup>H]-SK&F-107260 by 50% at equilibrium). The IC<sub>50</sub> is related to the equilibrium dissociation constant (K<sub>i</sub>) of the antagonist based on the Cheng and Prusoff equation:  $K_i = IC_{50} / (1 + L / K_d)$ , where L is the concentration of [<sup>3</sup>H]-SK&F-107260 used in the competitive binding assay (4.5 nM), and K<sub>d</sub> is the dissociation constant of [<sup>3</sup>H]-SK&F-107260 which is 4.5 nM as determined by Scatchard analysis. The compounds of this invention inhibit [<sup>3</sup>H]-SK&F 107260 binding with K<sub>i</sub> in the range of about 2 nM to about 1.0 µM. Preferred compounds generally have K<sub>i</sub> of less than 60 nM.

25

#### Inhibition of Platelet Aggregation

Blood was collected (citrate to prevent coagulation) from, naive, adult mongrel dogs. Platelet rich plasma, PRP, was prepared by centrifugation at 150 x g for 10 min at room temperature. Washed platelets were prepared by centrifuging PRP at 800 x g for 10 min. The cell pellet thus obtained was washed twice in Tyrode's buffer (pH 6.5) without Ca<sup>++</sup> and resuspended in Tyrode's buffer (pH 7.4) containing 1.8 mM Ca<sup>++</sup> at 3 x 10<sup>5</sup> cells/ml. Peptides were added 3 min prior to the agonist in all assays of platelet aggregation. Final agonist concentrations were 0.1 unit/ml thrombin and 2 mM ADP (Sigma). Aggregation was monitored in a Chrono-Log Lumi-Aggregometer. Light transmittance 5 min after addition of the agonist was used to calculate percent aggregation according to the formula % aggregation = [(90-CR) ÷ (90-10)] x 100, where CR is the chart reading, 90 is the

baseline, and 10 is the PRP blank reading. IC<sub>50</sub>'s were determined by plotting [% inhibition of aggregation] vs. [concentration of peptide]. Peptides were assayed at 200 mM and diluted sequentially by a factor of 2 to establish a suitable dose response curve.

- 5           The compounds of this invention inhibit the aggregation of human platelets stimulated with ADP with IC<sub>50</sub> of about 0.02 to about 2.0 μM. Preferred compounds have IC<sub>50</sub> of less than 1 μM. The most preferred compounds have IC<sub>50</sub> of less than 0.1 μM.

- 10           To assess the stability of the compounds to plasma proteases, the compounds were incubated for 3 h (rather than 3 min) in the PRP prior to addition of the agonist.

#### In Vivo Inhibition of Platelet Aggregation

- 15           *In vivo* inhibition of thrombus formation is demonstrated by recording the systemic and hemodynamic effects of infusion of the peptides into anesthetized dogs according to the methods described in Aiken *et al.*, *Prostaglandins*, 19, 629 (1980).

#### Inhibition of vitronectin binding

- 20           *Solid-Phase [<sup>3</sup>H]-SK&F-107260 Binding to α<sub>v</sub>β<sub>3</sub>*: Human placenta or human platelet α<sub>v</sub>β<sub>3</sub> (0.1-0.3 mg/mL) in buffer T (containing 2 mM CaCl<sub>2</sub> and 1% octylglucoside) was diluted with buffer T containing 1 mM CaCl<sub>2</sub>, 1 mM MnCl<sub>2</sub>, 1 mM MgCl<sub>2</sub> (buffer A) and 0.05% NaN<sub>3</sub>, and then immediately added to 96-well ELISA plates (Corning, New York, NY) at 0.1 mL per well. 0.1 - 0.2 μg of α<sub>v</sub>β<sub>3</sub>  
25           was added per well. The plates were incubated overnight at 4°C. At the time of the experiment, the wells were washed once with buffer A and were incubated with 0.1 mL of 3.5% bovine serum albumin in the same buffer for 1 hr at room temperature. Following incubation the wells were aspirated completely and washed twice with 0.2 mL buffer A.  
30           Compounds were dissolved in 100% DMSO to give a 2 mM stock solution, which was diluted with binding buffer (15 mM Tris-HCl (pH 7.4), 100 mM NaCl, 1 mM CaCl<sub>2</sub>, 1 mM MnCl<sub>2</sub>, 1 mM MgCl<sub>2</sub>) to a final compound concentration of 100 μM. This solution is then diluted to the required final compound concentration. Various concentrations of unlabeled antagonists (0.001 - 100 μM) were added to the  
35           wells in triplicates, followed by the addition of 5.0 nM of [<sup>3</sup>H]-SK&F-107260 (65 - 86 Ci/mmol).

The plates were incubated for 1 hr at room temperature. Following incubation the wells were aspirated completely and washed once with 0.2 mL of ice cold buffer A in a well-to-well fashion. The receptors were solubilized with 0.1 mL of 1% SDS and the bound [<sup>3</sup>H]-SK&F-107260 was determined by liquid  
5 scintillation counting with the addition of 3 mL Ready Safe in a Beckman LS Liquid Scintillation Counter, with 40% efficiency. Nonspecific binding of [<sup>3</sup>H]-SK&F-107260 was determined in the presence of 2 μM SK&F-107260 and was consistently less than 1% of total radioligand input. The IC<sub>50</sub> (concentration of the antagonist to inhibit 50% binding of [<sup>3</sup>H]-SK&F-107260) was determined by a  
10 nonlinear, least squares curve-fitting routine, which was modified from the LUNDON-2 program. The K<sub>i</sub> (dissociation constant of the antagonist) was calculated according to the equation:  $K_i = IC_{50} / (1 + L/K_d)$ , where L and K<sub>d</sub> were the concentration and the dissociation constant of [<sup>3</sup>H]-SK&F-107260, respectively.

Compounds of the present invention inhibit vitronectin binding to SK&F  
15 107260 in the concentration range of about 15 to greater than 100 micromolar. Preferred compounds inhibit vitronectin binding at a concentration of less than 50 micromolar.

Compounds of this invention are also tested for *in vitro* and *in vivo* bone resorption in assays standard in the art for evaluating inhibition of bone formation,  
20 such as the pit formation assay disclosed in EP 528 587, which may also be performed using human osteoclasts in place of rat osteoclasts, and the ovariectomized rat model, described by Wronski *et al.*, *Cells and Materials* 1991, Sup. 1, 69-74.

25 The examples which follow are intended to in no way limit the scope of this invention, but are provided to illustrate how to make and use the compounds of this invention. Many other embodiments will be readily apparent and available to those skilled in the art.

### 30 EXAMPLES

In the Examples, all temperatures are in degrees Centigrade. Mass spectra were performed using electro-spray (ES) ionization. Melting points were taken on a Thomas-Hoover capillary melting point apparatus and are uncorrected.

Celite® is filter aid composed of acid washed diatomaceous silica, and is a  
35 registered trademark of Mansville Corp., Denver, Colorado. Analtech silica gel GF and EM silica gel thin layer plates were used for thin layer chromatography. Both flash and gravity chromatography were carried out on Merck 60 (230-400 mesh)

silica gel. ODS refers to an octadecylsilyl derivatized silica gel chromatographic support.

Methyl (±)-7-carboxy-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate and methyl (±)-8-carboxy-3-oxo-4-(2-phenylethyl)-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate were prepared by the method of Bondinell, et al., WO 93/00095.

The following methods are illustrative of the manner of making certain useful intermediates for the preparation of the compounds of this invention.

10

#### Example 1

#### Preparation of (±)-7-[[bis[2-(4-pyridinyl)ethyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

## 15 a) Bis [2-(4-pyridyl)ethyl]amine

A mixture of 4-vinylpyridine (10.0 g, 95.1 mmole) and ammonium chloride (5.1 g, 95.1 mmole) in MeOH (90 mL) was heated to reflux under argon for 27 h. The resulting mixture was filtered and the filtrate was concentrated on the rotavap. The residue was dissolved in H<sub>2</sub>O (200 mL), and the solution was basified to pH 10.5 using 2 N NaOH. CH<sub>2</sub>Cl<sub>2</sub> extraction (3x100 mL), drying (MgSO<sub>4</sub>), and concentration gave a yellow oil (5.73 g, 49% crude). 2.9 g of this oil was chromatographed on silica gel (12% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to afford the title compound as a yellow oil (1.81 g, 16 %). MS (ES) m/e 228 (M + H)<sup>+</sup>.

## 25 b) Methyl (±)-7-[[bis[2-(4-pyridinyl)ethyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate

Diisopropylethylamine (0.35 g, 2.7 mmole) was added in one portion to a stirred mixture of bis [2-(4-pyridyl)ethyl]amine (0.37 g, 1.65 mmole), methyl (±)-7-carboxy-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (0.44 g, 1.50 mmole), EDC (0.34 g, 1.8 mmole), and HOBT · H<sub>2</sub>O (0.24 g, 1.8 mmole) in DMF (8 mL) at 0°C under argon, and the reaction was allowed to warm to RT. After 19.5 h, the reaction was poured into a mixture of ice water (100 g) and 5% NaHCO<sub>3</sub> (10 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub> (2x100 mL). The combined organic layers were washed with 5% NaHCO<sub>3</sub> (50 mL), dried (MgSO<sub>4</sub>), and concentrated. Chromatography on silica gel (7 % MeOH/CH<sub>2</sub>Cl<sub>2</sub>) gave the title compound (0.7 g, 88%). MS (ES) m/e 502.4 (M + H)<sup>+</sup>.

30

35



c) (±)-7-[[Bis[2-(4-pyridinyl)ethyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

1.0 N LiOH (0.66 mL, 0.66 mmole) was added dropwise to a mixture of methyl (±)-7-[[bis[2-(4-pyridinyl)ethyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (0.16 g, 0.3 mmole) in THF (7 mL) and H<sub>2</sub>O (10 mL) at RT. After 22.5 h, the reaction mixture was concentrated on the rotavap. The resulting residue was neutralized with 1.0 N AcOH (1.5 mL) and kept in the refrigerator to give a crystalline product (0.114 g, 73%). Purification by ODS bond elute chromatography (CH<sub>3</sub>CN/H<sub>2</sub>O) afforded the title compound. MS (ES) m/e 488.2 (M + H)<sup>+</sup>.

### Example 2

Preparation of (±)-7-[[bis[2-(4-piperidinyl)ethyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

a) Methyl (±)-7-[[bis[2-(4-piperidinyl)ethyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate

A mixture of methyl (±)-7-[[bis[2-(4-pyridinyl)ethyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (0.35 g, 0.65 mmole), PtO<sub>2</sub> (0.11 g, 0.48 mmole), 1.0 N HCl (1.3 mL, 1.3 mmole), and MeOH (30 mL) was hydrogenated at 45 psi in a Parr apparatus. After 5 h, the reaction mixture was filtered through celite® and concentrated to give the crude title compound (0.6 g) which was used without further purification. MS(ES) m/e 514.4 (M + H)<sup>+</sup>.

b) (±)-7-[[Bis[2-(4-piperidinyl)ethyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

Following the procedure of Example 1(c), except substituting methyl (±)-7-[[bis[2-(4-piperidinyl)ethyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (0.32 g, 0.51 mmole) for methyl (±)-7-[[bis[2-(4-pyridinyl)ethyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate, the title compound (0.41 g) was prepared. MS (ES) m/e 500.5 (M+H)<sup>+</sup>.

### Example 3

Preparation of (±)-8-[[bis[2-(4-pyridinyl)ethyl]amino]carbonyl]-3-oxo-4-(2-phenylethyl)-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

a) Methyl (±)-8-[[bis[2-(4-pyridinyl)ethyl]amino]carbonyl]-3-oxo-4-(2-phenylethyl)-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate

Following the procedure of Example 1(b), except substituting methyl (±)-8-carboxy-3-oxo-4-(2-phenylethyl)-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (0.57 g, 1.5 mmole) for methyl (±)-7-carboxy-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate, the title compound (0.83 g, 93%) was prepared. MS (ES) m/e 592.4 (M + H)<sup>+</sup>.

10 b) (±)-8-[[Bis[2-(4-pyridinyl)ethyl]amino]carbonyl]-3-oxo-4-(2-phenylethyl)-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

Following the procedure of Example 1(c), except substituting methyl (±)-8-[[bis[2-(4-pyridinyl)ethyl]amino]carbonyl]-3-oxo-4-(2-phenylethyl)-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (0.15 g, 0.25 mmole) for methyl (±)-7-  
15 [[bis[2-(4-pyridinyl)ethyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate, the title compound (0.21 g) was prepared. MS (ES) m/e 578.2 (M + H)<sup>+</sup>.

#### Example 4

20 Preparation of (±)-8-[[bis[2-(4-piperidinyl)ethyl]amino]carbonyl]-4-(2-cyclohexylethyl)-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

a) Methyl (±)-8-[[bis[2-(4-piperidinyl)ethyl]amino]carbonyl]-4-(2-cyclohexylethyl)-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate

Following the procedure of Example 2(a), except substituting methyl (±)-8-[[bis[2-(4-pyridinyl)ethyl]amino]carbonyl]-3-oxo-4-(2-phenylethyl)-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (0.53 g, 0.90 mmole) for methyl (±)-7-  
[[bis[2-(4-pyridinyl)ethyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate, the title compound (0.63 g) was prepared. MS (ES)  
30 m/e 305.6 (M + 2 H)<sup>2+</sup>.

b) (±)-8-[[Bis[2-(4-piperidinyl)ethyl]amino]carbonyl]-4-(2-cyclohexylethyl)-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

Following the procedure of Example 1(c), except substituting methyl (±)-8-  
35 [[bis[2-(4-piperidinyl)ethyl]amino]carbonyl]-4-(2-cyclohexylethyl)-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (0.26 g, 0.38 mmole) for methyl (±)-7-[[bis[2-(4-pyridinyl)ethyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-



1,4-benzodiazepine-2-acetate, the crude title compound was prepared. ODS chromatography (step gradient, 5 - 22 % CH<sub>3</sub>CN/H<sub>2</sub>O containing 0.1% TFA) afforded the title compound (0.25 g). MS (ES) m/e 596.4 (M + H)<sup>+</sup>.

5

Example 5Parenteral Dosage Unit Composition

A preparation which contains 20 mg of the compound of Example 1 as a sterile dry powder is prepared as follows: 20 mg of the compound is dissolved in 15 mL of distilled water. The solution is filtered under sterile conditions into a 25 mL multi-dose ampoule and lyophilized. The powder is reconstituted by addition of 20 mL of 5% dextrose in water (D5W) for intravenous or intramuscular injection. The dosage is thereby determined by the injection volume. Subsequent dilution may be made by addition of a metered volume of this dosage unit to another volume of D5W for injection, or a metered dose may be added to another mechanism for dispensing the drug, as in a bottle or bag for IV drip infusion or other injection-infusion system.

15

Example 6Oral Dosage Unit Composition

A capsule for oral administration is prepared by mixing and milling 50 mg of the compound of Example 1 with 75 mg of lactose and 5 mg of magnesium stearate. The resulting powder is screened and filled into a hard gelatin capsule.

20

Example 7Oral Dosage Unit Composition

A tablet for oral administration is prepared by mixing and granulating 20 mg of sucrose, 150 mg of calcium sulfate dihydrate and 50 mg of the compound of Example 1 with a 10% gelatin solution. The wet granules are screened, dried, mixed with 10 mg starch, 5 mg talc and 3 mg stearic acid; and compressed into a tablet.


30

The foregoing is illustrative of the making and using of this invention. This invention, however, is not limited to the precise embodiments described herein, but encompasses all modifications within the scope of the claims which follow.

35

What is claimed is:

1. A compound of the formula:



wherein:

A is NR', CHR', O or S;

X<sup>1</sup> and X<sup>2</sup> are C=O or CH<sub>2</sub>, with the proviso that only one of X<sup>1</sup> or X<sup>2</sup> is

10 C=O;

R<sup>1</sup> is H, C<sub>1-6</sub>alkyl or (CHR')<sub>m</sub>-Y;

R<sup>2</sup> is -(CH<sub>2</sub>)<sub>t</sub>-CO<sub>2</sub>R<sup>3</sup>;

R<sup>3</sup> is H, C<sub>1-6</sub>alkyl or (CHR')<sub>m</sub>-Ar;

R<sup>4</sup> and R<sup>4'</sup> independently are -(CH<sub>2</sub>)<sub>s</sub>-N or -(CH<sub>2</sub>)<sub>s</sub>-NR'R'

15 Y is H, Ar, C<sub>3-7</sub>cycloalkyl, CO<sub>2</sub>R<sup>3</sup> or Tet;

R' is H, C<sub>1-6</sub>alkyl or (CH<sub>2</sub>)<sub>m</sub>-Ar;

Ar is phenyl or naphthyl unsubstituted or substituted by one to three  
C<sub>1-6</sub>alkyl, trifluoromethyl, halogen, OR', SR', CONR'R<sup>1</sup>, CO<sub>2</sub>R<sup>1</sup>, NO<sub>2</sub> or R<sup>5</sup>R<sup>1</sup>N;

R<sup>5</sup> is H, C<sub>1-6</sub>alkyl, (CHR')<sub>m</sub>-Y or CO(CHR')<sub>m</sub>-Y;

20 N is pyrroline, pyrrolidine, imidazole, imidazoline, imidazolidine, pyrazole,  
pyrazoline, pyrazolidine, piperidine, piperazine, morpholine, pyridine, pyridinium,  
tetrahydropyridine, tetrahydroazepine or hexahydroazepine;

m is 0 to 6;

s is 1 to 4; and

25 t is 1 or 2;

or a pharmaceutically acceptable salt thereof.

2. The compound according to claim 1 wherein X<sup>1</sup> is CH<sub>2</sub> and X<sup>2</sup> is  
30 C=O.

3. The compound according to claim 2 wherein A is NR'.

4. The compound according to claim 3 wherein  $R^1$  is  $C_{1-4}$ alkyl or  $(CHR')_n-Y$ , in which  $R'$  is H or  $C_{1-4}$ alkyl,  $m$  is 1 to 3 and  $Y$  is Ar or  $C_{3-7}$ cycloalkyl.
5. The compound according to claim 4 wherein  $R^4$  and  $R^{4'}$  independently are  $-(CH_2)_s-\textcircled{N}$ , in which  $\textcircled{N}$  is piperidine, piperazine, pyridine, pyridinium or tetrahydropyridine and  $s$  is 1 to 3.
6. The compound according to claim 5 wherein  $\textcircled{N}$  is pyridine.
7. The compound according to claim 6 which is:  
( $\pm$ )-7-[[bis[2-(4-pyridinyl)ethyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid or  
( $\pm$ )-8-[[bis[2-(4-pyridinyl)ethyl]amino]carbonyl]-3-oxo-4-(2-phenylethyl)-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid.
8. The compound according to claim 5 wherein  $\textcircled{N}$  is piperidine.
9. The compound according to claim 8 which is:  
( $\pm$ )-7-[[bis[2-(4-piperidinyl)ethyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid or  
( $\pm$ )-8-[[bis[2-(4-piperidinyl)ethyl]amino]carbonyl]-4-(2-cyclohexylethyl)-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid.
10. A pharmaceutical composition comprising a compound according to claim 1 and a pharmaceutically acceptable carrier.
11. A method for effecting inhibition of platelet aggregation which comprises administering a compound according to claim 1.
12. A method for treating stroke or a transient ischemia attack or myocardial infarction which comprises administering a compound according to claim 1.
13. A method for promoting reperfusion of an artery or vein and inhibiting reocclusion which comprises administering a fibrinolytic agent and a compound according to claim 1.

14. A method of inhibiting a vitronectin receptor which comprises administering a compound according to claim 1.

5 15. A method of inhibiting bone resorption in a mammal which comprises administering a compound according to claim 1.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US95/10670

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C07D 243/14, 223/16, 267/14, 281/10; A61K 31/55

US CL : 514/211, 213, 221; 540/490, 519, 523

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/211, 213, 221; 540/490, 519, 523

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  
Please See Extra Sheet.

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US, A, 5,158,947 (TATSUOKA ET AL.) 27 OCTOBER 1992.	1-15

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

15 NOVEMBER 1995

Date of mailing of the international search report

08 DEC 1995

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

ROBERT T. BOND Jr.

Telephone No. (703) 308-1235

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US95/10670

## B. FIELDS SEARCHED

Documentation other than minimum documentation that are included in the fields searched:

CHEMICAL ABSTRACTS VOL. 51 (1951) VOL. 121 (DEC. 1994)

1,4 BENZODIAZEPINE-2-ACETIC ACID

1,4 BENZODIAZEPINE-3-ONE